

Cosmological constraints from X-ray AGN clustering and SNIa data

S. Basilakos¹ & M. Plionis^{1,2}.

¹ *Institute of Astronomy & Astrophysics, National Observatory of Athens, I. Metaxa & V. Pavlou, Palaia Penteli, 15236 Athens, Greece*

² *Instituto Nacional de Astrofísica, Óptica y Electrónica (INAOE) Apartado Postal 51 y 216, 72000, Puebla, Pue., Mexico*

2 February 2008

ABSTRACT

We put constraints on the main cosmological parameters of different spatially flat cosmological models by combining the recent clustering results of XMM-Newton soft (0.5–2 keV) X-ray sources, which have a redshift distribution with median redshift $z \sim 1.2$, and SNIa data. Using a likelihood procedure we find that the model which best reproduces the observational data and that is consistent with stellar ages is the *concordance* Λ CDM model with: $\Omega_m \simeq 0.28$, $w \simeq -1$, $H_0 \simeq 72 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $t_0 \simeq 13.5 \text{ Gyr}$ and has an X-ray AGN clustering evolution which is constant in physical coordinates. For a different clustering evolution model (constant in comoving coordinates) we find another viable model, although less probable due to the smaller age of the Universe, with $\Omega_m \simeq 0.38$, $w \simeq -1.25$, $H_0 \simeq 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$ and $t_0 \simeq 12.9 \text{ Gyr}$.

Key words: galaxies: clustering- X-ray sources - cosmology:theory - large-scale structure of universe

1 INTRODUCTION

Recent advances in observational cosmology, based on the analysis of a multitude of high quality observational data (Type Ia supernovae, cosmic microwave background (CMB), large scale structure, age of the globular clusters, high redshift galaxies), strongly indicated that we are living in a flat ($\Omega_{\text{tot}} = 1$) accelerating Universe containing a small baryonic component, non-baryonic cold dark matter (CDM) to explain the clustering of extragalactic sources and an extra component with negative pressure, usually named “dark energy”, to explain the present accelerated expansion of the universe (eg. Riess, et al. 1998; Perlmutter et al. 1999; Efsthathiou et al. 2002; Spergel et al. 2003; Percival et al. 2002; Tonry et al. 2003; Schuecker et al. 2003; Riess et al. 2004; Tegmark et al. 2004)

The last few years there have been many theoretical speculations regarding the nature of the exotic “dark energy”. Various candidates have been proposed in the literature, among which the time varying Λ -parameter (eg. Ozer & Taha 1987), a scalar field having a self-interaction potential $V(\Phi)$ with the field energy density decreasing with a slower rate than the matter energy density (dubbed also “quintessence”, eg. Peebles & Ratra 2003 and references therein) or an extra “matter” component, which is described by an equation of state $p_Q = w\rho_Q$ with $w < -1/3$ (a redshift dependence of w

is also possible but present measurements are not precise enough to allow meaningful constraints; eg. Dicus & Repko 2004). A particular case of “dark energy” is the traditional Λ -model which corresponds to $w = -1$. Note that a variety of observations indicate that $w < -0.6$ for a flat geometry (eg. Ettori, Tozzi & Rosati 2002; Tonry et al. 2003; Riess et al. 2004; Schuecker 2005).

In this paper we put constraints on spatially flat cosmological models using the recently derived clustering properties of the XMM-Newton soft (0.5–2 keV) X-ray point sources (Basilakos et al. 2005), the SNIa data (Tonry et al. 2003) and the age of globular clusters (eg. Krauss 2003; Cayrel et al. 2001). Hereafter will use the normalized Hubble constant $H_0 = 100 h \text{ km s}^{-1} \text{ Mpc}^{-1}$.

2 X-RAY AGN CLUSTERING

In a previous paper (Basilakos et al 2005) we derived the angular correlation function of the soft (0.5–2 keV) XMM X-ray sources using a shallow (2–10 ksec) wide-field survey ($\sim 2.3 \text{ deg}^2$). A full description of the data reduction, source detection and flux estimation are presented in Georgakakis et al. (2004). Note that the survey contains 432 point sources within an effective area of $\sim 2.1 \text{ deg}^2$ (for $f_x \geq 2.7 \times 10^{-14} \text{ erg cm}^{-2} \text{ s}^{-1}$), while for $f_x \geq 8.8 \times 10^{-15} \text{ erg cm}^{-2} \text{ s}^{-1}$ the effective area of the survey is $\sim 1.8 \text{ deg}^2$. In Basilakos et al (2005) we

present the details of the correlation function estimation, the various biases that should be taken into account (the amplification bias and integral constraint), the survey luminosity and selection functions as well as issues related to possible stellar contamination. In particular, the redshift selection function of our X-ray sources, derived by using the soft-band luminosity function of Miyaji, Hasinger & Schmidt (2000), and assuming the realistic luminosity dependent density evolution of our sources, predicts a characteristic depth of $z \simeq 1.2$ for our sample (for details see Basilakos et al. 2005).

Our aim here is to compare the theoretical clustering predictions from different flat cosmological models to the actual observed angular clustering of distant X-ray AGNs. For the purpose of this study we use Limber's formula which relates the angular, $w(\theta)$, and the spatial, $\xi(r)$, correlation functions. In the case of a spatially flat Universe, Limber's equation can be written as:

$$w(\theta) = 2 \frac{\int_0^\infty \int_0^\infty x^4 \phi^2(x) \xi(r, z) dx du}{[\int_0^\infty x^2 \phi(x) dx]^2}, \quad (1)$$

where $\phi(x)$ is the selection function (the probability that a source at a distance x is detected in the survey) and x is the coordinate distance related to the redshift through

$$x(z) = \frac{c}{H_0} \int_0^z \frac{dy}{E(y)}, \quad (2)$$

with $E(z) = [\Omega_m(1+z)^3 + (1-\Omega_m)(1+z)^\beta]^{1/2}$ and $\beta = 3(1+w)$. The number of objects within a shell $(z, z+dz)$ and in a given survey of solid angle ω_s is:

$$\frac{dN}{dz} = \omega_s x^2 n_s \phi(x) \left(\frac{c}{H_0} \right) E^{-1}(z). \quad (3)$$

where n_s is the comoving number density at zero redshift. Combining the above system of equations we obtain:

$$w(\theta) = 2 \frac{H_0}{c} \int_0^\infty \left(\frac{1}{N} \frac{dN}{dz} \right)^2 E(z) dz \int_0^\infty \xi(r, z) du \quad (4)$$

where r is the physical separation between two sources, having an angular separation, θ , given by $r \simeq (1+z)^{-1} (u^2 + x^2 \theta^2)^{1/2}$ (small angle approximation). Therefore, in order to estimate the expected $w(\theta)$ in a cosmological model we also need to determine the source redshift distribution dN/dz , which as we said previously, is estimated by integrating the appropriate Miyaji et al. (2000) luminosity function.

2.1 The Role and Evolution of Galaxy Bias

It has been claimed that the large scale clustering of different mass tracers (galaxies or clusters) is biased with respect to the matter distribution (cf. Kaiser 1984; Bardeen 1986). It is also an essential ingredient for cold dark matter (CDM) models to reproduce the observed galaxy distribution (cf. Davis et al. 1985). Within the framework of linear biasing (cf. Kaiser 1984; Benson et al. 2000), the mass-tracer and dark-matter spatial correlations, at some redshift z , are related by:

$$\xi(r, z) = \xi_{DM}(r, z) b^2(z), \quad (5)$$

where $b(z)$ is the bias evolution function. This has been shown to be a monotonically increasing function of redshift (Mo & White 1996; Matarrese et al. 1997; Basilakos & Plionis 2001 and references therein). Here we use the bias evolution model of Basilakos & Plionis (2001, 2003), which is based on linear perturbation theory and the Friedmann-Lemaître solutions of the cosmological field equations. We remind the reader that for the case of a spatially flat cosmological model our general bias evolution can be written as:

$$b(z) = \mathcal{A}E(z) + \mathcal{C}E(z) \int_z^\infty \frac{(1+y)^3}{E^3(y)} dy + 1. \quad (6)$$

Note that our model gives a family of bias curves, due to the fact that it has two unknowns (the integration constants $\mathcal{A}, \mathcal{C}$). In this paper, for simplicity, we fix the value of $\mathcal{C}$ to $\simeq 0.004$, as was determined in Basilakos & Plionis (2003) from the $2dF$ galaxy correlation function. We have tested the robustness of our results by increasing $\mathcal{C}$ by a factor of 10 and 100 to find differences of only $\sim 5\%$ in the fitted values of Ω_m and b_0 . This behaviour can be explained from the fact that the dominant term in the right hand side of eq. (6) is the first term $[\propto (1+z)^{3/2}]$ while the second term has a slower dependence on redshift $[\propto (1+z)]$.

2.2 Clustering Evolution

The redshift evolution of the spatial mass correlation function, $\xi_{DM}(r, z)$, can be written as the Fourier transform of the spatial power spectrum $P(k)$. Using also eq. (5) we have:

$$\xi(r, z) = \frac{(1+z)^{-(3+\epsilon)} b^2(z)}{2\pi^2} \int_0^\infty k^2 P(k) \frac{\sin(kr)}{kr} dk, \quad (7)$$

where k is the comoving wavenumber. Note that the parameter ϵ parametrizes the type of clustering evolution (eg. de Zotti et al. 1990). If $\epsilon = \gamma - 3$ (with γ the slope of the spatial correlation function; $\gamma = 1.8$) the clustering is constant in comoving coordinates while if $\epsilon = -3$ the clustering is constant in physical coordinates.

The power spectrum of our CDM models is given by $P(k) \propto k^n T^2(k)$ with scale-invariant ($n = 1$) primeval inflationary fluctuations and $T(k)$ the CDM transfer function. In particular, we use the transfer function parameterization as in Bardeen et al. (1986), with the corrections given approximately by Sugiyama (1995) while the normalization of the power spectrum is given by: $\sigma_8 \simeq 0.5 \Omega_m^{-\gamma}$ with $\gamma \simeq 0.21 - 0.22w + 0.33 \Omega_m$ (Wang & Steinhardt 1998). We caution that this fit, based on the rich cluster abundances, has been derived for $w \geq -1$. In this work we assume that the fit is valid also for $w < -1$. Note that we also use the non-linear corrections introduced by Peacock & Dodds (1994).

3 COSMOLOGICAL CONSTRAINTS

3.1 X-ray AGN Clustering likelihood

It has been shown that the application of the correlation function analysis on samples of high redshift galaxies

can be used as a useful tool for cosmological studies (eg. Matsubara 2004). In our case to constrain the cosmological parameters we utilize a standard χ^2 likelihood procedure to compare the measured XMM soft source angular correlation function (Basilakos et al. 2005) with the prediction of different spatially flat cosmological models. In particular, we define the likelihood estimator^{*} as: $\mathcal{L}^{\text{AGN}}(\mathbf{c}) \propto \exp[-\chi_{\text{AGN}}^2(\mathbf{c})/2]$ with:

$$\chi_{\text{AGN}}^2(\mathbf{c}) = \sum_{i=1}^n \left[\frac{w_{\text{th}}(\theta_i, \mathbf{c}) - w_{\text{obs}}(\theta_i)}{\sigma_i} \right]^2. \quad (8)$$

where $\mathbf{c}$ is a vector containing the cosmological parameters that we want to fit and σ_i the observed angular correlation function uncertainty. Here we work within the framework of a flat ($\Omega_{\text{tot}} = 1$) cosmology with primordial adiabatic fluctuations and baryonic density of $\Omega_b h^2 \simeq 0.022$ (eg. Kirkman et al. 2003). In this case the corresponding vector is $\mathbf{c} \equiv (\Omega_m, w, h, b_0)$. We sample the various parameters as follows: the matter density $\Omega_m \in [0.01, 1]$ in steps of 0.01; the equation of state parameter $w \in [-3, -0.35]$ in steps of 0.05, the dimensionless Hubble constant $h \in [0.5, 0.9]$ in steps of 0.02 and the X-ray sources bias at the present time $b_0 \in [0.5, 4]$ in steps of 0.05. Note that in order to investigate possible equations of state, we have allowed the parameter w to take values below -1. Such models correspond to the so called *phantom* cosmologies (eg. Caldwell 2002; Corasaniti et al. 2004).

The resulting best fit parameters, for the two clustering evolution models, are presented in the first two rows of Table 1. It is important to note that our estimates for the Hubble parameter h are in very good agreement with those derived ($h = 0.72 \pm 0.07$) by the HST key project (Freeman et al. 2001). In Fig.1 we present the 1σ , 2σ and 3σ confidence levels in the (w, h) and (w, b_0) planes by marginalizing the former over Ω_m and b_0 and the latter over Ω_m and h . It is evident that w is degenerate with respect to both h and the bias at the present time.

Figure 2 also shows the 1σ , 2σ and 3σ confidence levels (continuous lines) in the (Ω_m, w) plane by marginalizing over the Hubble constant and the bias factor at the present time[†]. The equation of state parameter likelihood is not constrained by this analysis and all the values in the interval $-3 \leq w \leq -0.35$ are acceptable within the 1σ uncertainty. Therefore, in order to put further constraints on w we additionally use the SNIa data (Tonry et al. 2003) as well as the so called age limit, given by the age of the oldest globular clusters in our Galaxy ($t_0 > 12.7$ Gyr; Krauss 2003; Cayrel et al. 2001 and references therein).

3.2 The likelihood from the SNIa

We use the sample of 172 supernovae of Tonry et al. (2003) in order to constrain Ω_m and the equation of




Figure 1. Likelihood contours in the (w, h) plane (left panel) and the (w, b_0) plane (right panel) for $\epsilon = -1.2$ (a similar degeneracy is true also for the $\epsilon = -3$ clustering evolution model). The contours are plotted where $-2\ln\mathcal{L}/\mathcal{L}_{\text{max}}$ is equal to 2.30, 6.16 and 11.83, corresponding to 1σ , 2σ and 3σ confidence level.

state in the framework of a flat geometry ($\Omega_{\text{tot}} = 1$). In this case, the corresponding vector $\mathbf{c}$ is: $\mathbf{c} \equiv (\Omega_m, w)$ and the likelihood function can be written as: $\mathcal{L}^{\text{SNIa}}(\mathbf{c}) \propto \exp[-\chi_{\text{SNIa}}^2(\mathbf{c})/2]$ with:

$$\chi_{\text{SNIa}}^2(\mathbf{c}) = \sum_{i=1}^{172} \left[\frac{\log D_L^{\text{th}}(z_i, \mathbf{c}) - \log D_L^{\text{obs}}(z_i)}{\sigma_i} \right]^2. \quad (9)$$

where $D_L(z)$ is the dimensionless luminosity distance, $D_L(z) = H_0(1+z)x(z)$ and z_i is the observed redshift. The green lines in Fig. 2 represents the 1σ , 2σ , and 3σ , confidence levels in the (Ω_m, w) plane. We find that the best fit solution is $\Omega_m = 0.30 \pm 0.04$ for $w > -1$, in complete agreement with previous SNIa studies (Tonry et al. 2003; Riess et al. 2004).

3.3 The joined likelihoods

In order to combine the X-ray AGN clustering properties with the SNIa data we perform a joined likelihood analysis and marginalizing the X-ray clustering results over h and b_0 , which are not constrained by the value of w (see Fig.1), we obtain: $\mathcal{L}^{\text{joint}}(\Omega_m, w) = \mathcal{L}^{\text{AGN}} \times \mathcal{L}^{\text{SNIa}}$. Taking also into account the age limit (the yellow area in Fig. 2) the likelihood for the $\epsilon = -1.2$ clustering evolution model peaks at $\Omega_m = 0.28 \pm 0.02$ with $w = -1.05^{+0.1}_{-0.2}$ (corresponding to $t_0 = 13.5$ Gyr) which is in excellent agreement with the WMAP results of Spergel et al. (2003) and the REFLEX X-ray clusters + SNIa results of Schuecker et al. (2003). For the $\epsilon = -3$ clustering evolution model we obtain $\Omega_m = 0.38 \pm 0.03$ with $w = -1.25^{+0.10}_{-0.25}$ (which corresponds to $t_0 = 12.9$ Gyr). The latter model appears to be marginally ruled out by the stellar ages. Note that the normalization of the power spectrum that corresponds to these models is $\sigma_8 \simeq 0.98$ and 0.90 , respectively.

It is evident that the combined likelihood analysis puts strong constraints on the value of w and once including stellar ages it appears to favor the standard *concordance* Λ CDM ($\Omega \simeq 0.3$, $w \simeq -1$) cosmological model as well as a comoving AGN clustering model ($\epsilon = -1.2$).

^{*} Likelihoods are normalized to their maximum values.

[†] Hereafter, when we marginalize over the Hubble constant we will use $h = 0.72$ for $\epsilon = -1.2$ and $h = 0.7$ for $\epsilon = -3$.

Table 1. Cosmological parameters from the likelihood analysis: The 1st column indicates the data used (the last two rows correspond to the joint likelihood analysis). Errors of the fitted parameters represent 1σ uncertainties. Note that for the joined analysis the corresponding results are marginalized over the Hubble constant and the bias factor at the present time, for which we use the values indicated.

Data	Ω_m	w	h	b_0	t_0	χ^2/dof
XMM ($\epsilon = -1.2$)	$0.31^{+0.16}_{-0.08}$	uncons. (w = -1)	$0.72^{+0.02}_{-0.18}$	$2.30^{+0.70}_{-0.20}$	13.0	0.82
XMM($\epsilon = -3.0$)	$0.38^{+0.02}_{-0.14}$	uncons. (w = -1)	$0.70^{+0.04}_{-0.16}$	$1.20^{+0.60}_{-0.30}$	12.6	0.84
XMM($\epsilon = -1.2$)/SNIa	0.28 ± 0.02	$-1.05^{+0.10}_{-0.20}$	0.72	2.30	13.5	0.87
XMM($\epsilon = -3.0$)/SNIa	0.38 ± 0.03	$-1.25^{+0.10}_{-0.25}$	0.70	1.20	12.9	0.85




Figure 2. Likelihood contours in the (Ω_m, w) plane. The contours correspond to 1σ (2.30), 2σ (6.16) and 3σ (11.83) confidence levels, using the two different clustering behaviors (left panel for $\epsilon = -3$ and right panel for $\epsilon = -1.2$). Note that the black and the green lines correspond to the X-ray AGN clustering and SNIa results, respectively while the yellow area is ruled out by the stellar ages.

However, the model with $w \simeq -1.25$, $\Omega_m \simeq 0.38$ and $\epsilon = -3$ cannot be ruled out at any significant level.

Many other recent analyses utilizing different combinations of data seem to agree with the former cosmological model. For example, Tegmark et al. (2004) used the WMAP CMB anisotropies in combination with the SDSS galaxy power spectrum and found a good Λ CDM fit with $\Omega_m = 0.30 \pm 0.04$ and $h = 0.70^{+0.04}_{-0.03}$ (see also Spergel et al. 2003; Percival et al. 2003; Schuecker et al. 2003). Also combining the gas fraction in relaxed X-ray luminous clusters with the CMB and SNIa has provided stringent constraints with $\Omega_m \simeq 0.3$ and $w \simeq -1$ (eg. Allen et al. 2004; Rapetti, Allen & Weller 2004).

4 CONCLUSIONS

We have combined for the first time the clustering properties of distant X-ray AGNs, identified as soft (0.5–2 keV) point sources in a shallow $\sim 2.3 \text{ deg}^2$ XMM survey, which have a z -distribution that peaks at $z \simeq 1.2$, with the SNIa data. From the X-ray AGN clustering likelihood analysis alone we constrain $h \simeq 0.72 \pm 0.03$ (where the uncertainty is found after marginalizing over w and b_0). From the joined likelihood analysis and taking into account stellar ages we constrain the matter density and the equation of state parameters. The best model appears to be one with $\Omega_m \simeq 0.28$, $w \simeq -1$ and

a stable in comoving coordinates X-ray AGN clustering model. However, the model with $\Omega_m \simeq 0.38$, $w \simeq -1.25$ and constant in physical coordinates ($\epsilon = -3$) X-ray AGN clustering, of which the predicted age is marginally consistent with stellar ages, cannot be excluded at any significant level by our analysis.

REFERENCES

- Allen, S. W., Schmidt, R. W., Ebeling, H., Fabian, A. C., Speybroeck, L., 2004, MNRAS, 353, 457
- Bardeen, J.M., Bond, J.R., Kaiser, N. & Szalay, A.S., 1986, ApJ, 304, 15
- Basilakos, S. & Plionis, M., 2001, ApJ, 550, 522
- Basilakos, S. & Plionis, M., 2003, ApJ, 593, L61
- Basilakos, S., Plionis, M., Georgantopoulos, I., Georgakakis, A., 2005, MNRAS, 356, 183
- Benson A. J., Cole S., Frenk S. C., Baugh M. C., & Lacey G. C., 2000, MNRAS, 311, 793
- Caldell, R. R., 2002, Physics Letters B, 545, 23
- Cayrel, R., et al., 2001, Nature, 409, 691
- Corasaniti, P. S., Kunz, M., Parkinson, D., Copeland, E. J., Bassett, B. A., 2004, Phys. Rev. Lett., 80, 3006
- Davis, M., Efstathiou, G., Frenk, C. S., White, S.D.M., 1985, ApJ, 292, 371
- Dicus, D.A. & Repko, W.W., 2004, Phys.Rev.D, 70, 3527,
- de Zotti, G., Persic, M., Franceschini, A., Danese, L., Palumbo, G. G. C., Boldt, E. A., Marshall, F. E., 1990, ApJ, 351, 22
- Efstathiou, G., 2002, MNRAS, 330, L29
- Ettori, S., Tozzi, P., Rosati, P., 2003, A&A, 398, 879
- Freedman, W., L., et al., 2001, ApJ, 553, 47
- Georgakakis, A., et al., 2004, MNRAS, 349, 135
- Kaiser N., 1984, ApJ, 284, L9
- Kirkman, D., Tytler, D., Suzuki, N., O’Meara, J.M., Lubin, D., 2003, ApJS, 149, 1
- Krauss, L.M., 2003, ApJ, 596, L1
- Matarrese S., Coles P., Lucchin F., Moscardini L., 1997, MNRAS, 286, 115
- Matsubara T., 2004, ApJ, 615, 573
- Miyaji, T., Hasinger, G., Schmidt, M., 2000, A&A, 353, 25
- Mo, H.J. & White, S.D.M 1996, MNRAS, 282, 347
- Ozer, M., & Taha, O., 1987, Nucl. Phys. B, 287, 776
- Peacock, A. J., & Dodds, S. J., 1994, MNRAS, 267, 1020
- Peebles P.J.E., Ratra, B., 2003, RvMP, 75, 559
- Perlmuter, S., et al., 1999, ApJ, 517, 565
- Persival, J.W., et al., 2002, MNRAS, 337, 1068
- Rapetti, D., Allen, W. S., Weller, J., MNRAS, in press, astro-ph/0409574
- Riess, A. G., et al., 1998, AJ, 116, 1009
- Riess, A. G., et al., 2004, ApJ, 607, 665
- Schuecker, P., 2005, astro-ph/0502234

- Schuecker, P., Caldwell, R.R., Bhinger, H., Collins, C.A.,
Guzzo, L., Weinberg, N.N., 2003, A&A, 402, 53
Spergel, D. N., et al., 2003, ApJs, 148, 175
Sugiyama, N., 1995, ApJS, 100, 281
Tegmark M., et al., 2004, Phys.Rev.D., 69, 3501
Tonry, et al. , 2003, ApJ, 594, 1
Wang, L. & Steinhardt, P.J., 1998, ApJ, 508, 483

Cosmological constraints from X-ray AGN clustering and SNIa data

S. Basilakos¹ & M. Plionis^{1,2}.

¹ *Institute of Astronomy & Astrophysics, National Observatory of Athens, I. Metaxa & V. Pavlou, Palaia Penteli, 15236 Athens, Greece*

² *Instituto Nacional de Astrofísica, Óptica y Electrónica (INAOE) Apartado Postal 51 y 216, 72000, Puebla, Pue., Mexico*

25 February 2005

ABSTRACT

We put constraints on the main cosmological parameters of different spatially flat cosmological models by combining the recent clustering results of XMM-*Newton* soft (0.5–2 keV) X-ray sources, which have a redshift distribution with median redshift $z \sim 1.2$, and SNIa data. Using a likelihood procedure we find that the model which best reproduces the observational data and that is consistent with stellar ages is the *concordance* Λ CDM model with: $\Omega_m \simeq 0.28$, $w \simeq -1$, $H_0 \simeq 72 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $t_0 \simeq 13.5 \text{ Gyr}$ and has an X-ray AGN clustering evolution which is constant in physical coordinates. For a different clustering evolution model (constant in comoving coordinates) we find another viable model, although less probable due to the smaller age of the Universe, with $\Omega_m \simeq 0.38$, $w \simeq -1.25$, $H_0 \simeq 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$ and $t_0 \simeq 12.9 \text{ Gyr}$.

Key words: galaxies: clustering- X-ray sources - cosmology:theory - large-scale structure of universe

1 INTRODUCTION

Recent advances in observational cosmology, based on the analysis of a multitude of high quality observational data (Type Ia supernovae, cosmic microwave background (CMB), large scale structure, age of the globular clusters, high redshift galaxies), strongly indicated that we are living in a flat ($\Omega_{\text{tot}} = 1$) accelerating Universe containing a small baryonic component, non-baryonic cold dark matter (CDM) to explain the clustering of extragalactic sources and an extra component with negative pressure, usually named “dark energy”, to explain the present accelerated expansion of the universe (eg. Riess, et al. 1998; Perlmutter et al. 1999; Efsthathiou et al. 2002; Spergel et al. 2003; Percival et al. 2002; Tonry et al. 2003; Schuecker et al. 2003; Riess et al. 2004; Tegmark et al. 2004)

The last few years there have been many theoretical speculations regarding the nature of the exotic “dark energy”. Various candidates have been proposed in the literature, among which the time varying Λ -parameter (eg. Ozer & Taha 1987), a scalar field having a self-interaction potential $V(\Phi)$ with the field energy density decreasing with a slower rate than the matter energy density (dubbed also “quintessence”, eg. Peebles & Ratra 2003 and references therein) or an extra “matter” component, which is described by an equation of state $p_Q = w\rho_Q$ with $w < -1/3$ (a redshift dependence of w

is also possible but present measurements are not precise enough to allow meaningful constraints; eg. Dicus & Repko 2004). A particular case of “dark energy” is the traditional Λ -model which corresponds to $w = -1$. Note that a variety of observations indicate that $w < -0.6$ for a flat geometry (eg. Ettori, Tozzi & Rosati 2002; Tonry et al. 2003; Riess et al. 2004; Schuecker 2005).

In this paper we put constraints on spatially flat cosmological models using the recently derived clustering properties of the XMM-*Newton* soft (0.5–2 keV) X-ray point sources (Basilakos et al. 2005), the SNIa data (Tonry et al. 2003) and the age of globular clusters (eg. Krauss 2003; Cayrel et al. 2001). Hereafter will use the normalized Hubble constant $H_0 = 100 h \text{ km s}^{-1} \text{ Mpc}^{-1}$.

2 X-RAY AGN CLUSTERING

In a previous paper (Basilakos et al 2005) we derived the angular correlation function of the soft (0.5–2 keV) XMM X-ray sources using a shallow (2–10 ksec) wide-field survey ($\sim 2.3 \text{ deg}^2$). A full description of the data reduction, source detection and flux estimation are presented in Georgakakis et al. (2004). Note that the survey contains 432 point sources within an effective area of $\sim 2.1 \text{ deg}^2$ (for $f_x \geq 2.7 \times 10^{-14} \text{ erg cm}^{-2} \text{ s}^{-1}$), while for $f_x \geq 8.8 \times 10^{-15} \text{ erg cm}^{-2} \text{ s}^{-1}$ the effective area of the survey is $\sim 1.8 \text{ deg}^2$. In Basilakos et al (2005) we

present the details of the correlation function estimation, the various biases that should be taken into account (the amplification bias and integral constraint), the survey luminosity and selection functions as well as issues related to possible stellar contamination. In particular, the redshift selection function of our X-ray sources, derived by using the soft-band luminosity function of Miyaji, Hasinger & Schmidt (2000), and assuming the realistic luminosity dependent density evolution of our sources, predicts a characteristic depth of $z \simeq 1.2$ for our sample (for details see Basilakos et al. 2005).

Our aim here is to compare the theoretical clustering predictions from different flat cosmological models to the actual observed angular clustering of distant X-ray AGNs. For the purpose of this study we use Limber's formula which relates the angular, $w(\theta)$, and the spatial, $\xi(r)$, correlation functions. In the case of a spatially flat Universe, Limber's equation can be written as:

$$w(\theta) = 2 \frac{\int_0^\infty \int_0^\infty x^4 \phi^2(x) \xi(r, z) dx du}{[\int_0^\infty x^2 \phi(x) dx]^2}, \quad (1)$$

where $\phi(x)$ is the selection function (the probability that a source at a distance x is detected in the survey) and x is the coordinate distance related to the redshift through

$$x(z) = \frac{c}{H_0} \int_0^z \frac{dy}{E(y)}, \quad (2)$$

with $E(z) = [\Omega_m(1+z)^3 + (1-\Omega_m)(1+z)^\beta]^{1/2}$ and $\beta = 3(1+w)$. The number of objects within a shell $(z, z+dz)$ and in a given survey of solid angle ω_s is:

$$\frac{dN}{dz} = \omega_s x^2 n_s \phi(x) \left(\frac{c}{H_0} \right) E^{-1}(z). \quad (3)$$

where n_s is the comoving number density at zero redshift. Combining the above system of equations we obtain:

$$w(\theta) = 2 \frac{H_0}{c} \int_0^\infty \left(\frac{1}{N} \frac{dN}{dz} \right)^2 E(z) dz \int_0^\infty \xi(r, z) du \quad (4)$$

where r is the physical separation between two sources, having an angular separation, θ , given by $r \simeq (1+z)^{-1} (u^2 + x^2 \theta^2)^{1/2}$ (small angle approximation). Therefore, in order to estimate the expected $w(\theta)$ in a cosmological model we also need to determine the source redshift distribution dN/dz , which as we said previously, is estimated by integrating the appropriate Miyaji et al. (2000) luminosity function.

2.1 The Role and Evolution of Galaxy Bias

It has been claimed that the large scale clustering of different mass tracers (galaxies or clusters) is biased with respect to the matter distribution (cf. Kaiser 1984; Bardeen 1986). It is also an essential ingredient for cold dark matter (CDM) models to reproduce the observed galaxy distribution (cf. Davis et al. 1985). Within the framework of linear biasing (cf. Kaiser 1984; Benson et al. 2000), the mass-tracer and dark-matter spatial correlations, at some redshift z , are related by:

$$\xi(r, z) = \xi_{DM}(r, z) b^2(z), \quad (5)$$

where $b(z)$ is the bias evolution function. This has been shown to be a monotonically increasing function of redshift (Mo & White 1996; Matarrese et al. 1997; Basilakos & Plionis 2001 and references therein). Here we use the bias evolution model of Basilakos & Plionis (2001, 2003), which is based on linear perturbation theory and the Friedmann-Lemaître solutions of the cosmological field equations. We remind the reader that for the case of a spatially flat cosmological model our general bias evolution can be written as:

$$b(z) = AE(z) + CE(z) \int_z^\infty \frac{(1+y)^3}{E^3(y)} dy + 1. \quad (6)$$

Note that our model gives a family of bias curves, due to the fact that it has two unknowns (the integration constants A, C). In this paper, for simplicity, we fix the value of C to $\simeq 0.004$, as was determined in Basilakos & Plionis (2003) from the $2dF$ galaxy correlation function. We have tested the robustness of our results by increasing C by a factor of 10 and 100 to find differences of only $\sim 5\%$ in the fitted values of Ω_m and b_0 . This behaviour can be explained from the fact that the dominant term in the right hand side of eq. (6) is the first term $[\propto (1+z)^{3/2}]$ while the second term has a slower dependence on redshift $[\propto (1+z)]$.

2.2 Clustering Evolution

The redshift evolution of the spatial mass correlation function, $\xi_{DM}(r, z)$, can be written as the Fourier transform of the spatial power spectrum $P(k)$. Using also eq. (5) we have:

$$\xi(r, z) = \frac{(1+z)^{-(3+\epsilon)} b^2(z)}{2\pi^2} \int_0^\infty k^2 P(k) \frac{\sin(kr)}{kr} dk, \quad (7)$$

where k is the comoving wavenumber. Note that the parameter ϵ parametrizes the type of clustering evolution (eg. de Zotti et al. 1990). If $\epsilon = \gamma - 3$ (with γ the slope of the spatial correlation function; $\gamma = 1.8$) the clustering is constant in comoving coordinates while if $\epsilon = -3$ the clustering is constant in physical coordinates.

The power spectrum of our CDM models is given by $P(k) \propto k^n T^2(k)$ with scale-invariant ($n = 1$) primeval inflationary fluctuations and $T(k)$ the CDM transfer function. In particular, we use the transfer function parameterization as in Bardeen et al. (1986), with the corrections given approximately by Sugiyama (1995) while the normalization of the power spectrum is given by: $\sigma_8 \simeq 0.5 \Omega_m^{-\gamma}$ with $\gamma \simeq 0.21 - 0.22w + 0.33 \Omega_m$ (Wang & Steinhardt 1998). We caution that this fit, based on the rich cluster abundances, has been derived for $w \geq -1$. In this work we assume that the fit is valid also for $w < -1$. Note that we also use the non-linear corrections introduced by Peacock & Dodds (1994).

3 COSMOLOGICAL CONSTRAINTS

3.1 X-ray AGN Clustering likelihood

It has been shown that the application of the correlation function analysis on samples of high redshift galaxies

can be used as a useful tool for cosmological studies (eg. Matsubara 2004). In our case to constrain the cosmological parameters we utilize a standard χ^2 likelihood procedure to compare the measured XMM soft source angular correlation function (Basilakos et al. 2005) with the prediction of different spatially flat cosmological models. In particular, we define the likelihood estimator^{*} as: $\mathcal{L}^{\text{AGN}}(\mathbf{c}) \propto \exp[-\chi^2_{\text{AGN}}(\mathbf{c})/2]$ with:

$$\chi^2_{\text{AGN}}(\mathbf{c}) = \sum_{i=1}^n \left[\frac{w_{\text{th}}(\theta_i, \mathbf{c}) - w_{\text{obs}}(\theta_i)}{\sigma_i} \right]^2. \quad (8)$$

where $\mathbf{c}$ is a vector containing the cosmological parameters that we want to fit and σ_i the observed angular correlation function uncertainty. Here we work within the framework of a flat ($\Omega_{\text{tot}} = 1$) cosmology with primordial adiabatic fluctuations and baryonic density of $\Omega_b h^2 \simeq 0.022$ (eg. Kirkman et al. 2003). In this case the corresponding vector is $\mathbf{c} \equiv (\Omega_m, w, h, b_0)$. We sample the various parameters as follows: the matter density $\Omega_m \in [0.01, 1]$ in steps of 0.01; the equation of state parameter $w \in [-3, -0.35]$ in steps of 0.05, the dimensionless Hubble constant $h \in [0.5, 0.9]$ in steps of 0.02 and the X-ray sources bias at the present time $b_0 \in [0.5, 4]$ in steps of 0.05. Note that in order to investigate possible equations of state, we have allowed the parameter w to take values below -1. Such models correspond to the so called *phantom* cosmologies (eg. Caldwell 2002; Corasaniti et al. 2004).

The resulting best fit parameters, for the two clustering evolution models, are presented in the first two rows of Table 1. It is important to note that our estimates for the Hubble parameter h are in very good agreement with those derived ($h = 0.72 \pm 0.07$) by the HST key project (Freeman et al. 2001). In Fig.1 we present the 1σ , 2σ and 3σ confidence levels in the (w, h) and (w, b_0) planes by marginalizing the former over Ω_m and b_0 and the latter over Ω_m and h . It is evident that w is degenerate with respect to both h and the bias at the present time.

Figure 2 also shows the 1σ , 2σ and 3σ confidence levels (continuous lines) in the (Ω_m, w) plane by marginalizing over the Hubble constant and the bias factor at the present time[†]. The equation of state parameter likelihood is not constrained by this analysis and all the values in the interval $-3 \leq w \leq -0.35$ are acceptable within the 1σ uncertainty. Therefore, in order to put further constraints on w we additionally use the SNIa data (Tonry et al. 2003) as well as the so called age limit, given by the age of the oldest globular clusters in our Galaxy ($t_0 > 12.7$ Gyr; Krauss 2003; Cayrel et al. 2001 and references therein).

3.2 The likelihood from the SNIa

We use the sample of 172 supernovae of Tonry et al. (2003) in order to constrain Ω_m and the equation of

^{*} Likelihoods are normalized to their maximum values.

[†] Hereafter, when we marginalize over the Hubble constant we will use $h = 0.72$ for $\epsilon = -1.2$ and $h = 0.7$ for $\epsilon = -3$.




Figure 1. Likelihood contours in the (w, h) plane (left panel) and the (w, b_0) plane (right panel) for $\epsilon = -1.2$ (a similar degeneracy is true also for the $\epsilon = -3$ clustering evolution model). The contours are plotted where $-2\ln\mathcal{L}/\mathcal{L}_{\text{max}}$ is equal to 2.30, 6.16 and 11.83, corresponding to 1σ , 2σ and 3σ confidence level.

state in the framework of a flat geometry ($\Omega_{\text{tot}} = 1$). In this case, the corresponding vector $\mathbf{c}$ is: $\mathbf{c} \equiv (\Omega_m, w)$ and the likelihood function can be written as: $\mathcal{L}^{\text{SNIa}}(\mathbf{c}) \propto \exp[-\chi^2_{\text{SNIa}}(\mathbf{c})/2]$ with:

$$\chi^2_{\text{SNIa}}(\mathbf{c}) = \sum_{i=1}^{172} \left[\frac{\log D_L^{\text{th}}(z_i, \mathbf{c}) - \log D_L^{\text{obs}}(z_i)}{\sigma_i} \right]^2. \quad (9)$$

where $D_L(z)$ is the dimensionless luminosity distance, $D_L(z) = H_0(1+z)x(z)$ and z_i is the observed redshift. The green lines in Fig. 2 represent the 1σ , 2σ , and 3σ confidence levels in the (Ω_m, w) plane. We find that the best fit solution is $\Omega_m = 0.30 \pm 0.04$ for $w > -1$, in complete agreement with previous SNIa studies (Tonry et al. 2003; Riess et al. 2004).

3.3 The joined likelihoods

In order to combine the X-ray AGN clustering properties with the SNIa data we perform a joined likelihood analysis and marginalizing the X-ray clustering results over h and b_0 , which are not constrained by the value of w (see Fig.1), we obtain: $\mathcal{L}^{\text{joint}}(\Omega_m, w) = \mathcal{L}^{\text{AGN}} \times \mathcal{L}^{\text{SNIa}}$. Taking also into account the age limit (the yellow area in Fig. 2) the likelihood for the $\epsilon = -1.2$ clustering evolution model peaks at $\Omega_m = 0.28 \pm 0.02$ with $w = -1.05^{+0.1}_{-0.2}$ (corresponding to $t_0 = 13.5$ Gyr) which is in excellent agreement with the WMAP results of Spergel et al. (2003) and the REFLEX X-ray clusters + SNIa results of Schuecker et al. (2003). For the $\epsilon = -3$ clustering evolution model we obtain $\Omega_m = 0.38 \pm 0.03$ with $w = -1.25^{+0.10}_{-0.25}$ (which corresponds to $t_0 = 12.9$ Gyr). The latter model appears to be marginally ruled out by the stellar ages. Note that the normalization of the power spectrum that corresponds to these models is $\sigma_8 \simeq 0.98$ and 0.90 , respectively.

It is evident that the combined likelihood analysis puts strong constraints on the value of w and once including stellar ages it appears to favor the standard *concordance* Λ CDM ($\Omega \simeq 0.3$, $w \simeq -1$) cosmological model as well as a comoving AGN clustering model ($\epsilon = -1.2$).

Table 1. Cosmological parameters from the likelihood analysis: The 1st column indicates the data used (the last two rows correspond to the joint likelihood analysis). Errors of the fitted parameters represent 1σ uncertainties. Note that for the joined analysis the corresponding results are marginalized over the Hubble constant and the bias factor at the present time, for which we use the values indicated.

Data	Ω_m	w	h	b_0	t_0	χ^2/dof
XMM ($\epsilon = -1.2$)	$0.31^{+0.16}_{-0.08}$	uncons. (w = -1)	$0.72^{+0.02}_{-0.18}$	$2.30^{+0.70}_{-0.20}$	13.0	0.82
XMM ($\epsilon = -3.0$)	$0.38^{+0.02}_{-0.14}$	uncons. (w = -1)	$0.70^{+0.04}_{-0.16}$	$1.20^{+0.60}_{-0.30}$	12.6	0.84
XMM ($\epsilon = -1.2$)/SNIa	0.28 ± 0.02	$-1.05^{+0.10}_{-0.20}$	0.72	2.30	13.5	0.87
XMM ($\epsilon = -3.0$)/SNIa	0.38 ± 0.03	$-1.25^{+0.10}_{-0.25}$	0.70	1.20	12.9	0.85




Figure 2. Likelihood contours in the (Ω_m, w) plane. The contours correspond to 1σ (2.30), 2σ (6.16) and 3σ (11.83) confidence levels, using the two different clustering behaviors (left panel for $\epsilon = -3$ and right panel for $\epsilon = -1.2$). Note that the black and the green lines correspond to the X-ray AGN clustering and SNIa results, respectively while the yellow area is ruled out by the stellar ages.

However, the model with $w \simeq -1.25$, $\Omega_m \simeq 0.38$ and $\epsilon = -3$ cannot be ruled out at any significant level.

Many other recent analyses utilizing different combinations of data seem to agree with the former cosmological model. For example, Tegmark et al. (2004) used the WMAP CMB anisotropies in combination with the SDSS galaxy power spectrum and found a good Λ CDM fit with $\Omega_m = 0.30 \pm 0.04$ and $h = 0.70^{+0.04}_{-0.03}$ (see also Spergel et al. 2003; Percival et al. 2003; Schuecker et al. 2003). Also combining the gas fraction in relaxed X-ray luminous clusters with the CMB and SNIa has provided stringent constraints with $\Omega_m \simeq 0.3$ and $w \simeq -1$ (eg. Allen et al. 2004; Rapetti, Allen & Weller 2004).

4 CONCLUSIONS

We have combined for the first time the clustering properties of distant X-ray AGNs, identified as soft (0.5–2 keV) point sources in a shallow $\sim 2.3 \text{ deg}^2$ XMM survey, which have a z -distribution that peaks at $z \simeq 1.2$, with the SNIa data. From the X-ray AGN clustering likelihood analysis alone we constrain $h \simeq 0.72 \pm 0.03$ (where the uncertainty is found after marginalizing over w and b_0). From the joined likelihood analysis and taking into account stellar ages we constrain the matter density and the equation of state parameters. The best model appears to be one with $\Omega_m \simeq 0.28$, $w \simeq -1$ and

a stable in comoving coordinates X-ray AGN clustering model. However, the model with $\Omega_m \simeq 0.38$, $w \simeq -1.25$ and constant in physical coordinates ($\epsilon = -3$) X-ray AGN clustering, of which the predicted age is marginally consistent with stellar ages, cannot be excluded at any significant level by our analysis.

REFERENCES

- Allen, S. W., Schmidt, R. W., Ebeling, H., Fabian, A. C., Speybroeck, L., 2004, MNRAS, 353, 457
- Bardeen, J.M., Bond, J.R., Kaiser, N. & Szalay, A.S., 1986, ApJ, 304, 15
- Basilakos, S. & Plionis, M., 2001, ApJ, 550, 522
- Basilakos, S. & Plionis, M., 2003, ApJ, 593, L61
- Basilakos, S., Plionis, M., Georgantopoulos, I., Georgakakis, A., 2005, MNRAS, 356, 183
- Benson A. J., Cole S., Frenk S. C., Baugh M. C., & Lacey G. C., 2000, MNRAS, 311, 793
- Caldell, R. R., 2002, Physics Letters B, 545, 23
- Cayrel, R., et al., 2001, Nature, 409, 691
- Corasaniti, P. S., Kunz, M., Parkinson, D., Copeland, E. J., Bassett, B. A., 2004, Phys. Rev. Lett., 80, 3006
- Davis, M., Efstathiou, G., Frenk, C. S., White, S.D.M., 1985, ApJ, 292, 371
- Dicus, D.A. & Repko, W.W., 2004, Phys.Rev.D, 70, 3527,
- de Zotti, G., Persic, M., Franceschini, A., Danese, L., Palumbo, G. G. C., Boldt, E. A., Marshall, F. E., 1990, ApJ, 351, 22
- Efstathiou, G., 2002, MNRAS, 330, L29
- Ettori, S., Tozzi, P., Rosati, P., 2003, A&A, 398, 879
- Freedman, W., L., et al., 2001, ApJ, 553, 47
- Georgakakis, A., et al., 2004, MNRAS, 349, 135
- Kaiser N., 1984, ApJ, 284, L9
- Kirkman, D., Tytler, D., Suzuki, N., O'Meara, J.M., Lubin, D., 2003, ApJS, 149, 1
- Krauss, L.M., 2003, ApJ, 596, L1
- Matarrese S., Coles P., Lucchin F., Moscardini L., 1997, MNRAS, 286, 115
- Matsubara T., 2004, ApJ, 615, 573
- Miyaji, T., Hasinger, G., Schmidt, M., 2000, A&A, 353, 25
- Mo, H.J., & White, S.D.M 1996, MNRAS, 282, 347
- Ozer, M., & Taha, O., 1987, Nucl. Phys. B, 287, 776
- Peacock, A. J., & Dodds, S. J., 1994, MNRAS, 267, 1020
- Peebles P.J.E., Ratra, B., 2003, RvMP, 75, 559
- Perlmutter, S., et al., 1999, ApJ, 517, 565
- Persival, J.W., et al., 2002, MNRAS, 337, 1068
- Rapetti, D., Allen, W. S., Weller, J., MNRAS, in press, astro-ph/0409574
- Riess, A. G., et al., 1998, AJ, 116, 1009
- Riess, A. G., et al., 2004, ApJ, 607, 665
- Schuecker, P., 2005, astro-ph/0502234

- Schuecker, P., Caldwell, R.R., Bhringer, H., Collins, C.A.,
Guzzo, L., Weinberg, N.N., 2003, A&A, 402, 53
Spergel, D. N., et al., 2003, ApJs, 148, 175
Sugiyama, N., 1995, ApJS, 100, 281
Tegmark M., et al., 2004, Phys.Rev.D., 69, 3501
Tonry, et al. , 2003, ApJ, 594, 1
Wang, L. & Steinhardt, P.J., 1998, ApJ, 508, 483